




High-Quality Draft Genome Sequences of Five *Xanthomonas arboricola* pv. *fragariae* Isolates

Michael Gétaz,^a Steve Baeyen,^b Jochen Blom,^c Martine Maes,^b Bart Cottyn,^b  Joël F. Pothier^a

^aEnvironmental Genomics and Systems Biology Research Group, Institute for Natural Resource Sciences, Zurich University of Applied Sciences (ZHAW), Wädenswil, Switzerland

^bPlant Sciences Unit—Plant Health, Flanders Research Institute for Agriculture, Fisheries and Food (ILVO), Merelbeke, Belgium

^cBioinformatics and Systems Biology, Justus-Liebig University Giessen, Giessen, Germany

ABSTRACT *Xanthomonas arboricola* pv. *fragariae* was described in 2001 as the causal agent of strawberry bacterial leaf blight. We report here the first draft whole-genome sequences of five *X. arboricola* pv. *fragariae* isolates from Italy and France.

Xanthomonas arboricola pv. *fragariae* was described in 2001 as the causal agent of strawberry bacterial leaf blight (1). The first symptoms related to this bacterium were observed in 1993 in strawberry cultivations in northern Italy (1). Afterward, the disease has only been reported in plantlets from Turkey in 2004 (2). The bacterium was under quarantine status in Europe starting in 2002 (3), but it was derestricted in 2007 (see https://www.eppo.int/MEETINGS/2007_meetings/phytomeasures.htm). This was partly due to the observation that the pathogen was sometimes coisolated with *Xanthomonas fragariae* (4, 5), which is the causal agent of bacterial angular leaf spot and is under quarantine status in Europe (6, 7). Also, the reported pathogenicity of *X. arboricola* pv. *fragariae* upon artificial inoculation on strawberry has often been ambiguous (4, 5, 8; M. Gétaz and J. F. Pothier, unpublished data). This could be related to the important heterogeneity among *X. arboricola* pv. *fragariae* strains, as observed in previous studies (4, 5, 9–11). Very recently, the pathogenicity of the pathotype strain was proved to produce typical symptoms in the strawberry cultivars Candonga, Sabrina, and Murano (12).

Until now, no whole-genome data for this bacterium were available in GenBank. In this study, whole-genome sequences of five *X. arboricola* pv. *fragariae* strains isolated from strawberry plants in Italy and France between 1986 and 1993 were obtained (Table 1).

Genomic DNA was extracted using the NucleoSpin tissue kit (Macherey-Nagel AG, Düren, Germany) following the manufacturer's protocol. Paired-end libraries constructed by the Nextera XT DNA library prep kit (Illumina, San Diego, CA) were sequenced on a MiSeq system (Illumina) using a 600-cycle MiSeq reagent kit v3 (Illumina). *De novo* assemblies were created using SeqMan NGen from the Lasergene genomics package version 12.1.0 (DNASar, Madison, WI). This was followed by contig reassembly using SeqMan Pro and read mapping using SeqMan NGen to check for inconsistencies.

The five genomes displayed an overall size between 4,692,498 and 4,906,785 bp, which is in the range previously observed with other *X. arboricola* genomes (13–17). The G+C contents in all the genomes were similar (65.75 to 65.95%). Genomes were annotated automatically using GenDB (18), and a total of 3,852 to 4,044 coding sequences (CDSs) were detected (Table 1). Using EDGAR 2.0 (19), a total of 3,523 CDSs were found to be shared between these five genomes.

Received 20 December 2017 Accepted 9 January 2018 Published 15 February 2018

Citation Gétaz M, Baeyen S, Blom J, Maes M, Cottyn B, Pothier JF. 2018. High-quality draft genome sequences of five *Xanthomonas arboricola* pv. *fragariae* isolates. Genome Announc 6:e01585-17. <https://doi.org/10.1128/genomeA.01585-17>.

Copyright © 2018 Gétaz et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Joël F. Pothier, joel.pothier@zhaw.ch.

TABLE 1 Draft whole-genome sequences of five *Xanthomonas arboricola* pv. *fragariae* strains submitted to the ENA database

Strain ^a	Synonym(s)	Yr, country of isolation	ENA accession no.	No. of contigs	Genome size (bp)	G+C content (%)	No. of CDSs
LMG 19145 ^b	CFBP 6771, PD 2780	1993, Italy	OEQL00000000	39	4,906,785	65.88	4,019
LMG 19144	CFBP 6770, PD 2696	1993, Italy	OEQF00000000	58	4,842,182	65.94	3,954
CFBP 6762	PD 2694	1993, Italy	OEQE00000000	38	4,889,284	65.75	4,044
LMG 19146	CFBP 3548, PD 3164	1986, France	OEQG00000000	39	4,884,039	65.75	4,020
CFBP 6773	NA ^c	NA	OEQD00000000	39	4,692,498	65.95	3,852

^aThe culture collections providing strains are abbreviated in the strain names as LMG (Collection of the Laboratorium voor Microbiologie en Microbiële Genetica, Ghent, Belgium), CFBP (Collection Française de Bactéries Associées aux Plantes, Beaucauzé, France), or PD (Culture Collection of Plant Pathogenic Bacteria, Wageningen, the Netherlands).

^bPathotype strain.

^cNA, not applicable.

Average nucleotide identities (ANIs) between these five genomes ranged from 96.57% to 97.69%, thus confirming the species designation but also suggesting some genome content heterogeneity.

The *X. arboricola* pv. *fragariae* draft genome sequences presented here add to the existing genomic information and further clarify the complexity of the species *X. arboricola*.

Accession number(s). The annotated draft whole-genome sequences of the five *X. arboricola* pv. *fragariae* isolates were deposited at ENA under the sequencing project number PRJEB23514. The accession numbers for the isolates are shown in Table 1.

ACKNOWLEDGMENTS

We are thankful to Marilena Palmisano for her assistance in the lab.

The EDGAR platform is funded by BMBF grant FKZ 031A533. This research project has received funding from the European Union's Seventh Framework Programme (FP7/2007-2013) for research, technological development, and demonstration under grant agreement 613678 (DROPSA). This article is based on work from European Cooperation in Science and Technology (COST) action CA16107 EuroXanth, supported by COST.

REFERENCES

- Janse JD, Rossi MP, Gorkink RFJ, Derks JHJ, Swings J, Janssens D, Scortichini M. 2001. Bacterial leaf blight of strawberry (*Fragaria (x) ananassa*) caused by a pathovar of *Xanthomonas arboricola*, not similar to *Xanthomonas fragariae* Kennedy & King. Description of the causal organism as *Xanthomonas arboricola* pv. *fragariae* (pv. nov., comb. nov.). *Plant Pathol* 50:653–665. <https://doi.org/10.1046/j.1365-3059.2001.00644.x>.
- Ustun N, Tjou-Tam-Sin NNA, Janse JD. 2007. First report on bacterial leaf blight of strawberry caused by *Xanthomonas arboricola* pv. *fragariae* Janse et al. in Turkey. *J Plant Pathol* 89:109–112.
- Anonymous. 2002. A new pathogen, *Xanthomonas arboricola* pv. *fragariae*, causing bacterial leaf blight of strawberry: addition to the EPPO Alert List. *OEPP/EPPO Reporting Serv* 2012(4):10–11.
- Scortichini M, Rossi MP. 2003. Genetic diversity of *Xanthomonas arboricola* pv. *fragariae* strains and comparison with some other *X. arboricola* pathovars using repetitive PCR genomic fingerprinting. *J Phytopathol* 151:113–119. <https://doi.org/10.1046/j.1439-0434.2003.00591.x>.
- Vandroemme J, Cottyn B, Pothier JF, Pflüger V, Duffy B, Maes M. 2013. *Xanthomonas arboricola* pv. *fragariae*: what's in a name? *Plant Pathol* 62:1123–1131. <https://doi.org/10.1111/ppa.12028>.
- Anonymous. 1986. *Xanthomonas fragariae* Kennedy & King. *EPPO Bull* 16: 17–20.
- European Union. 2000. Council Directive 2000/29/EC of 8 May 2000 on protective measures against the introduction into the community of organisms harmful to plants and plant products and their spread within the community. Council of the European Union, Brussels, Belgium.
- Merda D, Bonneau S, Guimbaud J-F, Durand K, Brin C, Boureau T, Lemaire C, Jacques M-A, Fischer-Le Saux M. 2016. Recombination-prone bacterial strains form a reservoir from which epidemic clones emerge in agroecosystems. *Environ Microbiol Rep* 8:572–581. <https://doi.org/10.1111/1758-2229.12397>.
- Hajri A, Pothier JF, Fischer-Le Saux M, Bonneau S, Poussier S, Boureau T, Duffy B, Manceau C. 2012. Type three effector gene distribution and sequence analysis provide new insights into the pathogenicity of plant-pathogenic *Xanthomonas arboricola*. *Appl Environ Microbiol* 78:371–384. <https://doi.org/10.1128/AEM.06119-11>.
- Cesbron S, Pothier J, Gironde S, Jacques MA, Manceau C. 2014. Development of multilocus variable-number tandem repeat analysis (MLVA) for *Xanthomonas arboricola* pathovars. *J Microbiol Methods* 100:84–90. <https://doi.org/10.1016/j.mimet.2014.02.017>.
- Fischer-Le Saux M, Bonneau S, Essakhi S, Manceau C, Jacques MA. 2015. Aggressive emerging pathovars of *Xanthomonas arboricola* represent widespread epidemic clones distinct from poorly pathogenic strains, as revealed by multilocus sequence typing. *Appl Environ Microbiol* 81: 4651–4668. <https://doi.org/10.1128/AEM.00050-15>.
- Ferrante P, Scortichini M. 2017. *Xanthomonas arboricola* pv. *fragariae*: a confirmation of the pathogenicity of the pathotype strain. *Eur J Plant Pathol* <https://doi.org/10.1007/s10658-017-1326-2>.
- Ibarra Caballero J, Zerillo MM, Snelling J, Boucher C, Tisserat N. 2013. Genome sequence of *Xanthomonas arboricola* pv. *corylina*, isolated from Turkish filbert in Colorado. *Genome Announc* 1:e00246-00213. <https://doi.org/10.1128/genomeA.00246-13>.
- Garita-Cambronero J, Sena-Vélez M, Palacio-Bielsa A, Cubero J. 2014. Draft genome sequence of *Xanthomonas arboricola* pv. *pruni*, strain Xap33, causal agent of bacterial spot disease on almond. *Genome Announc* 2:e00440-14. <https://doi.org/10.1128/genomeA.00440-14>.
- Ignatov AN, Kyrova EI, Vinogradova SV, Kamionskaya AM, Schaad NW, Luster DG. 2015. Draft genome sequence of *Xanthomonas arboricola* strain 3004, a causal agent of bacterial disease on barley. *Genome Announc* 3:e01572-14. <https://doi.org/10.1128/genomeA.01572-14>.
- Pereira UP, Gouran H, Nascimento R, Adaskaveg JE, Goulart LR, Dandekar AM. 2015. Complete genome sequence of *Xanthomonas arboricola* pv. *juglandis* 417, a copper-resistant strain isolated from

- Juglans regia* L. Genome Announc 3:e01126-15. <https://doi.org/10.1128/genomeA.01126-15>.
17. Garita-Cambronero J, Palacio-Bielsa A, López MM, Cubero J. 2016. Draft genome sequence for virulent and avirulent strains of *Xanthomonas arboricola* isolated from *Prunus* spp. in Spain. Stand Genomic Sci 11:12. <https://doi.org/10.1186/s40793-016-0132-3>.
 18. Meyer F, Goesmann A, McHardy AC, Bartels D, Bekel T, Clausen J, Kalinowski J, Linke B, Rupp O, Giegerich R, Pühler A. 2003. GenDB—an open source genome annotation system for prokaryote genomes. Nucleic Acids Res 31:2187–2195. <https://doi.org/10.1093/nar/gkg312>.
 19. Blom J, Kreis J, Spänig S, Juhre T, Bertelli C, Ernst C, Goesmann A. 2016. EDGAR 2.0: an enhanced software platform for comparative gene content analyses. Nucleic Acids Res 44:W22–W28. <https://doi.org/10.1093/nar/gkw255>.