



High-Quality Draft Genome Sequence of *Xanthomonas arboricola* pv. *juglandis* CPBF 1521, Isolated from Leaves of a Symptomatic Walnut Tree in Portugal without a Past of Phytosanitary Treatment

 Camila Fernandes,^{a,b,c} Jochen Blom,^d Joël F. Pothier,^e  Fernando Tavares^{a,c}

^aCIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório Associado, Microbial Diversity and Evolution Group, Universidade do Porto, Vairão, Portugal

^bUnidade Estratégica de Investigação e Serviços de Sistemas Agrários e Florestais e Sanidade Vegetal, Instituto Nacional de Investigação Agrária e Veterinária, Oeiras, Portugal

^cFaculdade de Ciências, Departamento de Biologia, Universidade do Porto, Porto, Portugal

^dBioinformatics and Systems Biology, Justus-Liebig-University Giessen, Giessen, Germany

^eEnvironmental Genomics and Systems Biology Research Group, Institute for Natural Resource Sciences, Zurich University of Applied Sciences (ZHAW), Wädenswil, Switzerland

ABSTRACT Here, we report the draft genome sequence of *Xanthomonas arboricola* pv. *juglandis* CPBF 1521, isolated from symptomatic leaves of an ornamental walnut in a public site in Portugal without any record of phytosanitary treatment. This isolate may constitute a genomic reference of a wild-type strain in comparative genomics studies.

Xanthomonas arboricola pv. *juglandis* (*Gammaproteobacteria* class, *Xanthomonadales* order, *Xanthomonadaceae* family) is a threatening and important pathogen of the principal commercial nut trees Persian walnut and English walnut (*Juglans regia* L.) (1, 2). Diseases caused by *X. arboricola* pv. *juglandis* have been demonstrated by the development of several symptoms, namely the presence of necrotic lesions on leaves and fruits, the presence of external apical necrosis near the blossom end evolving into fruit necrosis, and the presence of vertical cankers, brown to black exudates, and distortions on trunks (3–5). Not surprisingly, *X. arboricola* pv. *juglandis* is responsible for increasing losses in walnut production resulting in a negative economic impact for walnut crop regions in many countries worldwide (1, 2, 6).

The present announcement reports the whole-genome sequence of a *X. arboricola* pv. *juglandis* strain, CPBF 1521, isolated in October 2014 from the leaves of an ornamental *J. regia* specimen in a public site in Loures, Portugal, showing typical symptoms of walnut bacterial blight, and for which no phytosanitary treatments were applied. This set of features suggests that this strain has not been exposed to selective pressures caused by phytosanitary treatments, such as copper-based compound sprays, making this genomic data set particularly interesting for comparative genomics studies.

X. arboricola pv. *juglandis* CPBF 1521 was obtained from infected leaf samples as previously described (7) and was grown on M2 medium (yeast extract, 2 g liter⁻¹; Bacto peptone, 5 g liter⁻¹; NaCl, 5 g liter⁻¹; KH₂PO₄, 0.45 g liter⁻¹; and Na₂HPO₄ 12H₂O, 2.39 g liter⁻¹) at 28°C for 48 h with shaking (100 rpm). The EZNA bacterial DNA purification kit (Omega Bio-Tek, Norcross, GA) was used for DNA extraction. Standard genomic library preparation and sequencing was carried out with at the GATC Biotech AG (Konstanz, Germany) using an Illumina HiSeq platform with 2 × 150-bp paired-end reads. Raw sequence data with approximately 10,113,730 reads were assembled

Received 10 August 2018 Accepted 2 October 2018 Published 25 October 2018

Citation Fernandes C, Blom J, Pothier JF, Tavares F. 2018. High-quality draft genome sequence of *Xanthomonas arboricola* pv. *juglandis* CPBF 1521, isolated from leaves of a symptomatic walnut tree in Portugal without a past of phytosanitary treatment. *Microbiol Resour Announc* 7:e00887-18. <https://doi.org/10.1128/MRA.00887-18>.

Editor Jason Stajich, University of California, Riverside

Copyright © 2018 Fernandes et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Camila Fernandes, camila.fernandes@cibio.up.pt, or Fernando Tavares, ftavares@fc.up.pt.

de novo using MIRA version 4.0 (8) in accurate mode with standard settings. The set of contigs obtained was reassembled using SeqMan Pro from the Lasergene genomics package version 12.1.0 (DNASTar, Madison, WI) with Pro Assembler parameters and reads were mapped using SeqMan NGen with standard settings to check for inconsistencies. Contigs were ordered according to the related reference genome of *X. arboricola* pv. *juglandis* (strain CFBP 2528; GenBank accession number [NZ_JZEF00000000](https://doi.org/10.1093/nar/nzjzef0000000)) (9) using the Move Contigs function in Mauve version 20150226 build 10 (10, 11). Genome annotation was performed with Prokka version 1.12 (12) based on *de novo* discovery of genes using a *Xanthomonas* genus database.

The *X. arboricola* pv. *juglandis* CPBF 1521 genome contains 56 contigs with an N_{50} value of 173,159 bp, has a total size of 5,194,740 bp, and a G+C content of 65.41%. A total of 4,494 coding sequences (CDS) with 4 rRNAs and 53 tRNAs were found using Prokka. Initial comparative analysis using EDGAR version 2.0 (13) identified 3,984 CDS shared between CPBF 1521 and *X. arboricola* pv. *juglandis* CFBP 2528 that was used as a reference genome. Moreover, a high average nucleotide identity value (98%) was obtained between these two strains.

This newly sequenced genome expands the existing genomic data for *X. arboricola* pv. *juglandis* (8, 14–16). Further genomic comparison studies are being conducted to identify unique genomic features of *X. arboricola* pv. *juglandis* CPBF 1521.

Data availability. This whole-genome shotgun project has been deposited in the European Nucleotide Archive (ENA) under the BioProject accession number [PRJEB27248](https://doi.org/10.6019/PRJEB27248) and in GenBank under the accession number [UIHD00000000](https://doi.org/10.1093/nar/uihd0000000). The version described in this paper is the first version, UIHD01000000.

ACKNOWLEDGMENTS

This work was cofinanced by the European Structural and Investment Funds (ESIFs) through the Operational Competitiveness and Internationalization Program, COMPETE 2020, and by national funds through Fundação para a Ciência e a Tecnologia (FCT), within the framework of the project EVOXANT (reference numbers PTDC/BIA-EVF/3635/2014 and POCI-01-0145-FEDER-016600). This article is based on work from European Cooperation in Science and Technology (COST) action CA16107 EuroXanth, supported by COST. Camila Fernandes was supported by a fellowship (SFRH/BD/95913/2013) from FCT and a short-term scientific mission (STSM) from COST action CA16107. The EDGAR platform is funded by BMBF grant number FKZ 031A533.

We thank Leonor Cruz for lab support.

REFERENCES

- Frutos D. 2010. Bacterial diseases of walnut and hazelnut and genetic resources. *J Plant Pathol* 92:579–585.
- Lamichhane JR. 2014. *Xanthomonas arboricola* diseases of stone fruit, almond, and walnut trees: progress toward understanding and management. *Plant Dis* 98:1600–1610. <https://doi.org/10.1094/PDIS-08-14-0831-FE>.
- Smith RE, Smith CO, Ramsey HJ. 1912. Walnut culture in California: walnut blight. California Agricultural Experiments Station Publications, Berkeley, CA.
- Moragrega C, Ozaktan H. 2010. Apical necrosis of Persian (English) walnut (*Juglans regia*): an update. *J Plant Pathol* 92:567–571.
- Hajri A, Meyer D, Delort F, Guillaumes J, Brin C, Manceau C. 2010. Identification of a genetic lineage within *Xanthomonas arboricola* pv. *juglandis* as the causal agent of vertical oozing canker of Persian (English) walnut in France. *Plant Pathol* 59:1014–1022. <https://doi.org/10.1111/j.1365-3059.2010.02362.x>.
- Moragrega C, Matias J, Aleta N, Montesinos E, Rovira M. 2011. Apical necrosis and premature drop of Persian (English) walnut fruit caused by *Xanthomonas arboricola* pv. *juglandis*. *Plant Dis* 95:1565–1570. <https://doi.org/10.1094/PDIS-03-11-0259>.
- Fernandes C, Albuquerque P, Sousa R, Cruz L, Tavares F. 2017. Multiple DNA markers for identification of *Xanthomonas arboricola* pv. *juglandis* isolates and its direct detection in plant samples. *Plant Dis* 101:858–865. <https://doi.org/10.1094/PDIS-10-16-1481-RE>.
- Chevreux B, Wetter T, Suhai S. 1999. Genome sequence assembly using trace signals and additional sequence information, p 45–56. In Wingender E (ed), *Computer science and biology. Proceedings of the German Conference on Bioinformatics, GCB 99, Hannover, Germany*.
- Cesbron S, Briand M, Essakhi S, Gironde S, Boureau T, Manceau C, Fischer-Le Saux M, Jacques MA. 2015. Comparative genomics of pathogenic and nonpathogenic strains of *Xanthomonas arboricola* unveil molecular and evolutionary events linked to pathoadaptation. *Front Plant Sci* 6:1126. <https://doi.org/10.3389/fpls.2015.01126>.
- Darling AC, Mau B, Blattner FR, Perna NT. 2004. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res* 14:1394–1403. <https://doi.org/10.1101/gr.2289704>.
- Darling AE, Mau B, Perna NT. 2010. progressiveMauve: multiple genome alignment with gene gain, loss and rearrangement. *PLoS One* 5:e11147. <https://doi.org/10.1371/journal.pone.0011147>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- Blom J, Kreis J, Spänig S, Juhre T, Bertelli C, Ernst C, Goesmann A. 2016. EDGAR 2.0: an enhanced software platform for comparative gene con-

- tent analyses. *Nucleic Acids Res* 44:W22–W28. <https://doi.org/10.1093/nar/gkw255>.
14. Higuera G, González-Escalona N, Véliz C, Vera F, Romero J. 2015. Draft genome sequences of four *Xanthomonas arboricola* pv. *juglandis* strains associated with walnut blight in Chile. *Genome Announc* 3:e01160-15. <https://doi.org/10.1128/genomeA.01160-15>.
 15. Pereira UP, Gouran H, Nascimento R, Adaskaveg JE, Goulart LR, Dandekar AM. 2015. Complete genome sequence of *Xanthomonas arboricola* pv. *juglandis* 417, a copper-resistant strain isolated from *Juglans regia* L. *Genome Announc* 3:e01126–e01115. <https://doi.org/10.1128/genomeA.01126-15>.
 16. Fu B, Chen Q, Wei M, Zhu J, Zou L, Li G, Wang L. 2018. Complete genome sequence of *Xanthomonas arboricola* pv. *juglandis* strain DW3F3, isolated from a *Juglans regia* L. bacterial blighted fruitlet. *Genome Announc* 6:e00023-18. <https://doi.org/10.1128/genomeA.00023-18>.